



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/82, A01H 5/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 97/48814</b> <b>(43) International Publication Date:</b> 24 December 1997 (24.12.97)
<b>(21) International Application Number:</b> PCT/US97/10621 <b>(22) International Filing Date:</b> 20 June 1997 (20.06.97) <b>(30) Priority Data:</b> 08/667,188 21 June 1996 (21.06.96) US <b>(71) Applicant:</b> MONSANTO COMPANY [US/US], 800 North Lindbergh Boulevard, St. Louis, MO 63167 (US). <b>(72) Inventors:</b> CHENG, Ming; 11532 Tivoli Lane, St. Louis, MO 63146 (US). FRY, Joyce, E.; 9839 Gerald Drive, St. Louis, MO 63128 (US). WAN, Yuechun; 893-F Foxsprings, Chesterfield, MO 63017 (US). <b>(74) Agent:</b> KITCHELL, Barbara, S.; Arnold, White & Durkee, P.O. Box 4433, Houston, TX 77210 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> METHODS FOR THE PRODUCTION OF STABLY-TRANSFORMED, FERTILE WHEAT EMPLOYING <i>AGROBACTERIUM</i> -MEDIATED TRANSFORMATION AND COMPOSITIONS DERIVED THEREFROM		
<b>(57) Abstract</b> <p>Disclosed are processes for producing stably transformed fertile wheat and a system of transforming wheat via <i>Agrobacterium</i>. This invention provides methods for transforming a variety of explants, such as freshly isolated or pre-cultured immature embryos, embryogenic callus and suspension cells. Also disclosed are methods for recovering transgenic plants after transformation within a short period of time, if the explants are regenerable at the time of transformation. Thus, the frequency of somaclonal variation associated with prolonged <i>in vitro</i> culture period is significantly reduced. The transformation frequency using this system is comparable to or better than published methods using other systems, such as microprojectile bombardment.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**METHODS FOR THE PRODUCTION OF STABLY-  
TRANSFORMED, FERTILE WHEAT EMPLOYING  
*AGROBACTERIUM*-MEDIATED TRANSFORMATION AND  
COMPOSITIONS DERIVED THEREFROM**

## 1. BACKGROUND OF THE INVENTION

### 1.1 Field of the Invention

The present invention relates to the field of molecular biology. More specifically, it concerns methods for the incorporation of foreign DNA into the genome of monocotyledonous plants, and in particular, wheat. Provided herein are reproducible systems for genetically transforming wheat, methods of selecting stable genetic transformants from suspensions of transformed cells, and methods of producing fertile plants from the transformed cells. Exemplary methods include the use of *Agrobacterium*-mediated transformation to introduce nucleic acids into cells, and selectable and/or screenable marker systems, for example, genes which confer resistance (*e.g.*, antibiotic, herbicide, *etc.*), or which contain an otherwise phenotypically observable trait. In other aspects, the invention relates to the production of stably transformed and fertile wheat plants, gametes and offspring from these plants.

### 1.2 Description of the Related Art

The entire text of U.S. patent application SN 08/329,742 filed October 26, 1994 is hereby incorporated by reference in its entirety. During the past decade, it has become possible to transfer genes from a wide range of organisms to crop plants by recombinant DNA technology. This advance has provided enormous opportunities to improve plant resistance to pests, diseases and herbicides, and to modify biosynthetic processes to change the quality of plant products (Knutson *et al.*, 1992; Piorer *et al.*, 1992; Vasil *et al.*, 1992). However, the availability of an efficient transformation method to introduce foreign DNA has been a substantial barrier for most monocot species, including maize, rice, oat, barley, and particularly wheat.

#### 1.2.1 Available Methods for Transforming Monocotyledonous Plants

There have been many methods attempted for the transformation of monocotyledonous plants but only a few methods have resulted in stable transformation. Two methods are currently employed for most transgenic studies in monocot species:

direct DNA transfer into isolated protoplasts and microprojectile-mediated DNA delivery (Shimamoto *et al.*, 1989; Fromm *et al.*, 1990). More recently, additional methods have also been developed for use in monocots. Following is a brief description of the methods that have resulted in stably transformed and fertile monocots capable of transferring genes to their progeny in a Mendelian fashion.

#### 1.2.1.1 Biolistics

"Biolistics" is most widely used transformation method for monocotyledons. In the "biolistics" method microprojectile particles are coated with DNA and accelerated by a mechanical device to a speed high enough to penetrate the plant cell wall and nucleus (Intl. Pat. Appl. Publ. No. WO 91/02071). The foreign DNA gets incorporated into the host DNA and results in a transformed cell. There are many variations on the "biolistics" method (Sanford, 1990; Fromm *et al.*, 1990; Christou *et al.*, 1988; Sautter *et al.*, 1991). This method has been successfully used to produce stably transformed monocotyledonous plants including rice, maize, wheat, barley, and oats (Christou *et al.*, 1991; Gordon-Kamm *et al.*, 1990; Vasil *et al.*, 1992, 1993; Wan *et al.*, 1993; Sommers *et al.*, 1992).

The microprojectile-mediated DNA delivery method may use immature embryos or immature embryo derived calli as target tissues. Transgenic plants have been recovered from the microprojectile bombardment method in maize, oat, barley and wheat (Gordon-Kamm *et al.*, 1990; Somers *et al.*, 1992; Wan *et al.*, 1994; Vasil *et al.*, 1992).

The microprojectile bombardment method generally takes 10 to 15 months to obtain transgenic plants (Gordon-Kamm *et al.*, 1990; Vasil *et al.*, 1992). Even with the more recent improvements in transformation methods using immature embryos as target tissues, it still requires 4 to 6 months to recover transgenic plants (Weeks *et al.*, 1993; Vasil *et al.*, 1992; 1993; Becker *et al.*, 1994). The transformation frequency by these methods is variable ranging from about one event from 100 to 1000 bombarded embryos.

### 1.2.1.2 Electroporation

The protoplast methods have been widely used in rice, where DNA is delivered to the protoplasts through liposomes, PEG, and electroporation. While a large number of transgenic plants have been recovered in several laboratories (Shimamoto *et al.*, 1989; Datta *et al.*, 1990), the protoplast methods require the establishment of long-term embryogenic suspension cultures. Some regenerants from protoplasts are infertile and phenotypically abnormal due to the long-term suspension culture (Davey *et al.*, 1991; Rhodes *et al.*, 1988). These procedures have been especially useful for rice and some grasses.

Transformation by electroporation involves the application of short, high voltage electric fields to create "pores" in the cell membrane through which DNA is taken-up. This method has been used to produce stably transformed monocotyledon plants, (Pasazkowski *et al.*, 1985; Shillito *et al.*, 1985; Fromm *et al.*, 1986) especially from rice (Shimamoto *et al.*, 1992; Datta *et al.*, 1990, 1992; Hayakawa *et al.*, 1992).

### 1.2.1.3 Chemical Treatment of Protoplasts

The polyethylene glycol (PEG) method is simply a chemical treatment in the presence of the protoplasts and the DNA (Shillito *et al.*, 1985; Rhodes *et al.*, 1988). The PEG facilitate the uptake of the DNA.

### 1.2.1.4 Other Methods

A number of other methods have been reported for the transformation of monocotyledon plants. The methods reported to produce fertile transgenic monocotyledon plants include the "pollen tube method" (Intl. Pat. Appl. Publ. No. WO 93/18168; Zahir, 1993, Luo and Wu, 1988) and macro-injection of DNA into floral tillers (Du *et al.*, 1989; Picard *et al.*, 1988; De la Pena *et al.*, 1987) and tissue incubation of seeds in DNA solutions (Tofer *et al.*, 1989). Direct injection of exogenous DNA into the endosperm of a fertilized plant ovule at the onset of embryogenesis was disclosed in Intl. Pat. Appl. Publ. No. WO 94/00583. Besides the protoplast and the biolistics methods of transformation other

methods are not reproducible or predictable. There is usually evidence of expression but seldom is the DNA transmitted to the progeny.

### 1.3 Deficiencies in the Prior Art

5           The one important area where there has been little significant progress in the art has been the adaptation of bacterial-mediated methods of transformation in monocots. While widely useful in dicotyledonous plants, *Agrobacterium*-mediated gene transfer has been disappointing when adapted to use in monocots. There are several reports in the literature claiming *Agrobacterium* transformation of monocotyledons which are discussed in Intl. Pat.  
10   Appl. Publ. No. WO 94/0077. These are specifically the methods of Gould *et al.*, 1991; Mooney *et al.*, 1991; and Raineri *et al.*, 1990, which claim *Agrobacterium* transformation of maize, rice and wheat. There is some evidence of gene transfer in these methods but they lack convincing evidence for transfer efficiency, reproducibility, and confirmation of gene transfer (Potrykus, 1990), and lack of transfer to the progeny when plants are produced. In  
15   the work of Gould where evidence of transformed plants was presented there was no Mendelian inheritance of the genes.

          De LaFonteyne *et al.* (Intl. Pat. Appl. Publ. No. WO 92/06205) described a process for the transformation of maize cells using *A. tumefaciens* strains in combination with a transposon-mediated integration method, but the success of such methods in other species  
20   was not demonstrated.

          Mooney *et al.* (1991) produced transformed cells from wheat embryos co-cultivated with *A. tumefaciens* but the frequency of transformation was very low and often unreproducible. Chan *et al.* (1993) subsequently attempted to produce transgenic rice plants using *A. tumefaciens*, but their methods have not been widely accepted owing  
25   to a lack of sufficient molecular and genetic evidence of transgenic plant production.

          More recent attempts by Hiei *et al.* (1994) suggested that transgenic rice plants could be obtained following *A. tumefaciens* transformation, but that the particular bacterial strains used and the choice of bacterial vectors were critical for successfully obtaining transgenes. A recent paper by Ishida *et al.* (1996) indicated a high-efficiency

transformation of maize was possible by co-culture of immature embryos with *A. tumefaciens*. In both reports on rice and the maize transformation, a super binary vector pTOK233 containing the *virB*, *virC* and *virG* genes was used to achieve high-efficiency transformation. A recent report by Saito *et al.* (Intl. Pat. Appl. Publ. No. WO 95/06722) discloses a method of transforming monocotyledons using scutellum of non-dedifferentiated immature embryos using *A. tumefaciens*.

Despite the fact that wheat is the most widely-grown cereal crop in the world, unfortunately no convincing reports exist on the use of *Agrobacterium* transformation methods in the preparation of stable, fertile wheat transgenic plants. Likewise, no methods have been developed using immature embryonic or callus tissues for stable, high-frequency transformation of wheat. Therefore, what is lacking in the prior art is an *Agrobacterium*-mediated method for preparing fertile, transgenic wheat plants.

## 2. SUMMARY OF THE INVENTION

The present invention seeks to overcome these and other inherent deficiencies in the prior art by providing novel processes for the stable transformation of monocotyledonous plants, and in particular, wheat, using *Agrobacterium*-mediated methods.

It is therefore a particular object of the present invention to provide techniques that will allow one to prepare transgenic, fertile wheat which are preferably diploid and which have been stably transformed through the introduction of one or more desired genes into the genome of these species.

The present invention thus relates generally to methods for the production of transgenic wheat plants. As used herein, the term transgenic plants is intended to refer to plants that have incorporated exogenous genes or DNA sequences, including but not limited to genes or DNA sequences which are perhaps not normally present, genes not normally transcribed and translated ("expressed") in a given cell type, or any other genes or DNA sequences which one desires to introduce into the non-transformed plant, such as genes which may normally be present in the non-transformed plant but which one desires



to have altered expression.

The present invention can be used with any plant species. It is particularly useful for monocot species. More particularly, it is useful in plant species which cannot remain in a callus state for long periods of time without losing the ability to regenerate. One particularly useful species in the present invention is wheat. Preferred species include *Triticum aestivum*, *T. turgidum* and *T. monococum* wheat, with *T. aestivum* being particularly preferred. The present invention, when applied to wheat, has the advantage of being genotype independent. That is, it can be used with any type of wheat variety, including both winter and spring wheat. It can be used to produce transgenic wheat plants from spring cultivars, such as, for example, Bobwhite and Marshall PAVOT1, UC702, and Panewawa as well as winter cultivars, such as, for example, HY368, Neeley, FL302, RH91, R332, R1269 and R585.

The present invention is used to introduce foreign DNA into regenerable plant tissue. Any type of foreign DNA can be inserted into the plant species using the method of the present invention. Generally, "foreign DNA" can be defined to include any type of DNA which is inserted into a plant cell from outside the plant cell. Methods for inserting cloned DNA into suitable plasmid constructs, and manipulation of appropriate *Agrobacterium* delivery strains are generally well known.

The type of DNA included in the foreign DNA can include DNA which already is present in the plant cell, DNA from another plant, DNA from a different organism, or a DNA generated externally, such as a DNA sequence containing an anti-sense message of a plant gene, or a DNA sequence encoding a synthetic version of a gene where the nucleotide sequence has been modified.

In a preferred embodiment, *Agrobacterium tumefaciens* C58, anopaline strain is used to mediate the transfer of DNA into a wheat cell. Other preferred strains for use in the practice of the invention include octopine strains, LBA4404 or agro pine strains, e.g., EHA101 or EHA105.

In preferred embodiments, the foreign DNA contains a DNA sequence which can function in a regenerable plant tissue as a selection device. Such DNA can include a gene

which would function in a regenerable plant tissue to produce a compound which would confer upon the plant tissue resistance to an otherwise toxic compound. These genes are well known in the art and can confer resistance to compounds such as antibiotics like kanamycin (Dekeyser *et al.*, 1989), and herbicides like glyphosate (Della-Cioppa *et al.*, 1987 and bialaphos (Vasil *et al.*, 1992). Other selection devices can be used within the scope of the present invention. Such genes include those for CP4, BAR, hygromycin, phosphotransferase (NPT) and dihydrofolate reductase (*dhfr*).

In one embodiment, a method is disclosed for producing a transgenic wheat plant.

The method involves in a general sense, establishing a culture from a wheat plant to be transformed, transforming the culture with an *Agrobacterium* comprising a DNA composition that includes a genetic component one desires to introduce into the wheat genome, identifying or selecting a transformed cell line, and regenerating a transgenic wheat plant therefrom.

In another important embodiment, the invention provides a method for producing a fertile transgenic wheat plant. The process involves establishing a regenerable culture from a wheat plant to be transformed, introducing a DNA composition comprising a genetic component one desires to introduce into the genome of said wheat plant, by *Agrobacterium* transformation, identifying or selecting a transformed cell line; and regenerating a fertile transgenic wheat plant from the transformed cell line. The DNA is transmitted through a complete sexual cycle of the transgenic plant to its progeny, and the progeny contain a stable, chromosomally-integrated copy of the selectable or screenable marker gene which was transformed into the parent via *Agrobacterium* transformation.

In these embodiments, the DNA composition comprises a plasmid, and particularly a recombinant plasmid such as pMON18365 which contains an *np11* gene. Other genes of interest include selectable or screenable marker genes such as *e.g.*, any of those described herein, including GUS, green fluorescent protein (GFP), luciferase (LUX), CP4 and *np11* genes. Examples of transposons and associated antibiotic resistance genes include the transposons Tns (*bla*), Tn5 (*np11*), Tn7 (*dhfr*), penicillins, kanamycin (and neomycin, G418, bleomycin); methotrexate (and trimethoprim);

chloramphenicol; kanamycin and tetracycline.

Characteristics useful for selectable markers in plants have been outlined in a report on the use of microorganisms (Advisory Committee on Novel Foods and Processes, July 1994). These include i) stringent selection with minimum number of  
5 nontransformed tissues; ii) large numbers of independent transformation events with no significant interference with the regeneration; iii) application to a large number of species; and iv) availability of an assay to score the tissues for presence of the marker. As mentioned, several antibiotic resistance markers satisfy these criteria, including those resistant to kanamycin (*npI*II) hygromycin B (*aph*IV) and gentamicin (*aac*C3 and *aac*C4).  
10 A more complete description and list is included in Table 1.

TABLE 1

Type	Examples
Aminoglycoside antibiotics	
1) phosphotransferase enzymes (APH)	APH (3')II APHIV
2) adenylyltransferase enzymes (AAD)	ADD (3")
3) acetyltransferases (AAC)	AAC(3)-I AAC(3)-III AAC(3)-IV
Chloramphenicol	
chloramphenicol acetyl transferase (CAT)	
$\beta$ -Lactam antibiotics	TEM-1- $\beta$ -lactamase
$\beta$ -lactamase	
2,4-Diaminopteridones dihydrofolate reductase	
Glycopeptides	TN5-bleomycin
Pyridone carboxylic acids	nalidixic acid resistance
DNA gyrase	
Rifamycins	rifamycin resistance
resistant RNA polymerase	
Macrolides	erythromycin resistance
50S subunit methylation	
Tetracyclines	excretes antibiotic from cell
bind to S4 and S18 proteins of 70S ribosome	
subunit	

Alternatives to antibiotic resistance markers have been developed for plants (Advisory Committee on Novel Foods and Processes, July 1994). These include metal tolerance markers, complementation systems with auxotrophic markers, sugar catabolism markers, L-canavanine resistance markers and markers for resistance to lysine and

threonine and S-aminoethyl L-cysteine. Examples and additional description are shown in Table 2.

TABLE 2

Marker	Example
Herbicide resistance	herbicide resistance can be used as a selectable marker in plants <i>e.g.</i> , tolerance to glyphosate and bromoynil
Metal tolerance	metal tolerance, through insertion of a mammalian metallothionein gene
Resistance to lysine and threonine and to s-aminoethyl L-cysteine	two of the enzymes of the aspartate family biosynthetic pathway, which is regulated by several feedback inhibition loops, have been developed as selectable markers; aspartate kinase (AK) activity is inhibited by millimolar concentrations of lysine and threonine (LT); dihydrodipicoline synthase (DHPS) activity is inhibited by lysine and leads to sensitivity to the toxic lysine analogue, S-aminoethyl L-cysteine (AEC); transgenic plants containing <i>E. coli</i> genes for expression of AK and DHPS can be grown on media containing LT and AEC respectively.

5

Scorable markers may also be used in plants as alternatives to antibiotic resistance markers. Such markers are used to detect the presence or to measure the level of expression of the transferred gene. The use of scorable markers in plants to identify or tag genetically modified cells works well only when efficiency of modification of the cell

is high. Some of the more commonly employed scorable markers include beta-glucuronidase (GUS), whose expression is detected by a blue color on incubation of the tissue with 5-bromo-4-chloro-3-indolyl-1-glucuronide; bacterial luciferase (LUX) whose expression is detected by light emission; firefly luciferase (LUC) whose  
5 expression is detected by light emission after incubation with luciferin; and  $\beta$ -galactosidase whose expression is detected by a bright blue color after the tissue is stained with 5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactopyranoside.

Preferred cells for practice of these methods include immature embryos, callus tissues, or suspension cells, and a preferred *Agrobacterium* species for transformation is  
10 *A. tumefaciens*, with strains such as C58 LBA4404 and EHA101 being particularly preferred.

The embryos to be transformed may either be freshly isolated from an immature caryopsis or isolated from an immature caryopsis and then pre-treated. The isolation of the immature embryo from the immature caryopsis may occur on the same day of  
15 inoculation, or alternatively, one, two, five, or even more days prior to inoculation. Likewise, the callus tissue may be isolated from an immature embryo or from non-embryonic plant cells. The embryo may be injured prior to transformation. The embryo or caryopsis may be one that is 2 or more days after anthesis, and when suspension cells are used, they may be cultures of individual cells or, alternatively, cell clusters.

20 The DNA to be transformed is preferably a recombinant plasmid, as described herein, and may comprise one or more promoters and/or 3' and/or 5' regions operatively linked to the particular genetic components to be transformed. Promoters such as CaMV 35S, FMV, ubiquitin, rice actin or enhanced CaMV 35S promoters are particularly preferred.

25 When desirable, the plants derived from the transformation may be grown and progeny obtained. These progeny may be used to prepare transgenic seeds, or alternatively, bred with a second transgenic wheat plant to prepare a fertile, transgenic wheat plant that comprises one or more transgenes of interest. The seeds obtained from such progeny may be germinated, cultivated, and used to prepare subsequent generations

of transgenic offspring which comprise the transgene originally transformed in the parental line.

In one embodiment of the present invention, an immature embryo from a plant is used as a starting material. Immature embryos can be produced using known method  
5 described in the art. For instance, the production of wheat immature embryos is described by Weeks *et al.* (1993) and Vasil *et al.* (1993).

In another preferred embodiment of the present invention, the regenerable plant tissues are calli. The preferred calli are embryogenic calli. Embryogenic calli are produced from immature embryos. These calli can be produced by isolating and  
10 culturing immature embryos on a nutrient media with carbohydrate and plant growth regulators. In the preferred embodiment of the present invention, when producing embryogenic calli from wheat, the elimination of embryo axis as described by Nehra *et al.*, (1994) is not necessary.

Callus-producing medium are well known in the art and any culture medium or  
15 preparation method can be used. In the preferred embodiment, where wheat calli are prepared, a wheat immature embryo is cultured for 1 day up to one month, preferably for 4 to 7 days, on a modified MS medium comprising about 40 g/l maltose and about 2 mg/l 2,4-D. In another embodiment, the 2,4-D can be replaced by a combination of 0.5 mg/l 2,4-D and 2.2 mg/l picloram (Chemservice). The medium is solidified by 2 g/l Gelrite®  
20 (Sigma Chemical, St. Louis, MO) or 4 g/l low-melting agarose.

After transformation, the regenerable plant tissue is placed in a medium capable of producing shoots from the regenerable tissue where the medium further contains a compound used to select regenerable tissue containing the selectable DNA sequences. This is in contrast to the prior art where regenerable plant tissue is generally subjected  
25 first to an extended period of selection prior to exposure of the regenerable tissue to a medium capable for producing shoots.

The medium used in this step can be any medium which permits the formation of shoots from the regenerable tissue. In one embodiment, a shoot-producing compound is added to the medium. These shoot-producing compounds are well known in the art

(Mursahige and Skoog, 1962; Kasha *et al.*, 1990). Such compounds include weak plant growth regulators and include IAA, IBA, and BA at low concentrations (Becker *et al.*, 1994; Vasil *et al.*, 1992). In another embodiment of the invention, a medium free of a plant growth regulator can be used to induce shoot formation (Weeks *et al.*, 1993).

- 5 In a preferred embodiment, where an embryogenic wheat callus is to be regenerated, the medium comprises a modified MS medium with 0.2 mg/l 2,4-D (Murashige and Skoog, 1962; Wan and Lemaux, 1994).

The regenerable plant tissue is generally placed in this medium as quickly as possible in the present invention after transformation. Generally, this can range from  
10 about 1 day to about three weeks, but preferably from about 1 day to about two weeks, more preferably from about two to about three weeks. Most preferably the tissue is transferred to this medium from about one week to about two weeks after transformation. In most instance, the transfer will occur between about 5 and about 11 days.

- 15 The compound used to select regenerable tissue containing the selectable DNA sequences can be any of a variety of well known selection compounds, such as antibiotics and herbicides. Preferred compounds can include geneticin (G-418) (aminoglycoside) (Nehra *et al.*, 1994), glyphosate (Della-Cioppa *et al.* 1987) and bialaphos (Vasil *et al.*, 1992; Weeks *et al.*, 1993).

- 20 The availability of alternative selection agents is an important requirement for commercial application of agriculture biotechnology. The use of kanamycin has been less successful for cereal crops because of the high endogenous level of tolerance (Dekeyser *et al.*, 1989). Bialaphos has been widely used as a selection agent in cereal crop transformation (Weeks *et al.*, 1993; Vasil *et al.*, 1993; Becker *et al.*, 1994; Nehra *et al.*, 1994; Wan and Lemaux, 1994). However, it could potentially be a disaster to  
25 exclusively use genes encoding bialaphos resistance as a selectable marker in all transformation studies. Other selectable markers are needed and the results demonstrate that the herein described rapid regeneration system works well with different selection agents.

After shoots have formed the shoots are transferred to a second medium capable



of producing roots from said shoots. This medium can further contain a compound used to select regenerable tissue containing the selectable DNA sequences. Transfer to this medium occurs when sufficient shoots have developed, a generally known in the art. This occurs, for wheat, within 25 to 40 days after transformation.

- 5           The medium capable of producing roots can be any root-producing medium. These mediums are well known in the art (Weeks *et al.*, 1993; Vasil *et al.*, 1992). One preferred root-producing medium is a modified MS medium without any plant growth regulator (Murashige and Skoog, 1962; Zhou *et al.*, 1992).

- 10           Once roots have been formed, the plants can then be transferred to soil and grown following methods known in the art to produce seeds.

- 15           The present invention discloses a reproducible efficient *Agrobacterium* method for transforming monocotyledons, especially wheat and corn with a conventional binary vector commonly used for transformation of dicotyledons. The invention provides a rapid and efficient *Agrobacterium* transformation and regeneration system, especially useful for the transformation of wheat and corn. Plants regenerated from this system are phenotypically normal and fully fertile. The transgenes are transmitted to R1 progeny in a Mendelian fashion.

- 20           In a preferred embodiment, the present invention provides a rapid and efficient transformation system for wheat using freshly isolated immature embryos, pre-cultured embryos and proliferated calli from immature embryos. The new transformation system takes about three months to obtain transgenic plants, and transformation frequencies with the novel methods disclosed herein are reproducibly 0.3 to 4.3%.

- 25           In a general sense, the present invention provides a method for producing a transformed monocotyledonous plant which contains exogenous DNA. Such a method generally involves isolating regenerable tissue from the plant, transferring into the regenerable tissue the foreign DNA by a 1 to 3 hour inoculation of the regenerable tissue with the *Agrobacterium* and co-culture of the plant tissue with the *Agrobacterium* for 2-3 days. Typically the foreign DNA to be inserted into the plant genome comprises a selectable marker DNA sequence, where the sequence can function in a regenerable tissue

as a selection device: The cells are then grown from between about 2 day to about 5 days on callus medium containing antibiotics to kill the *Agrobacterium*, and approximately 1 to 2 weeks later, the regenerable tissue is placed in a medium capable of producing shoots from the tissue. This medium further contains a compound used to select regenerable tissue containing the selectable DNA sequences; and after at least one shoot has formed, the shoot is typically transferred to a second medium capable of producing roots from the shoot.

## 2.2 Exogenous Genes

An aspect of the invention relates generally to transgenic plants which express one or more exogenous genes transformed via *A. tumefaciens*. As used herein, the term "transgenic plants" is intended to refer to plants that have incorporated DNA sequences, including but not limited to genes which are perhaps not normally present, DNA sequences not normally transcribed into RNA or translated into a protein ("expressed"), or any other genes or DNA sequences which one desires to introduce into the non-transformed plant, such as genes which may normally be present in the non-transformed plant but which one desires to either genetically engineer or to have altered expression. It is contemplated that in some instances the genome of transgenic plants of the present invention will have been augmented through the stable introduction of the transgene. However, in other instances, the introduced gene will replace an endogenous sequence.

Exemplary genes which may be introduced include, for example, DNA sequences or genes from another species, or even genes or sequences which originate with or are present in the same species, but are incorporated into recipient cells by genetic engineering methods rather than classical reproduction or breeding techniques. However, the term exogenous, is also intended to refer to genes which are not normally present in the cell being transformed, or perhaps simply not present in the form, structure, etc., as found in the transforming DNA segment or gene, or genes which are normally present yet which one desires, e.g., to have over-expressed. Thus, the term "exogenous" gene or DNA is intended to refer to any gene or DNA segment that is introduced into a recipient

cell, regardless of whether a similar gene may already be present in such a cell.

An initial step in the production of fertile transgenic plants is the obtaining of a DNA composition, *e.g.*, vectors, plasmids, linear DNA fragments, and the like, a component of which is to be delivered to recipient monocotyledonous cells. DNA segments for use in transforming such cells will, of course, generally comprise the gene or genes which one desires to introduce into the cells. These genes can further include structures such as promoters, enhancers, polylinkers, or even regulatory genes as desired.

Vectors, plasmids, cosmids, YACs (yeast artificial chromosomes) and DNA segments for use in transforming such cells will, of course, generally comprise the either cDNA, gene or gene sequences which one desires to transform into the monocotyledonous plants. These DNA constructs can further include structures such as promoters, enhancers, polylinkers, or even regulatory genes as desired. The DNA segment or gene may encode either a native or modified protein or polypeptide, which will be expressed in the resultant recombinant cells, and/or which will impart an improved phenotype to the regenerated plant.

In order to improve the ability to identify transformants, one may desire to employ a selectable or screenable marker gene as, or in addition to, the expressible gene of interest. Marker genes code for phenotypes that allow cells which express the marker gene to be distinguished from cells that do not have the marker. Such genes may encode either a selectable or screenable marker, depending on whether the marker confers a trait which one can select for by chemical means, *i.e.*, through the use of a selective agent (*e.g.*, a herbicide, antibiotic, or the like). Many examples of suitable marker genes are known in the art and may be employed in the practice of the invention.

Selectable markers for use in connection with the present invention include but are not limited to an *nrpII* gene; a *bar* gene which codes for bialaphos resistance; a mutant EPSP synthase gene which encodes glyphosate resistance; a nitrilase gene which confers resistance to bromoxynil; a mutant acetolactate synthase gene (ALS) which confers imidazolinone or sulphonylurea resistance; or a methotrexate resistant DHFR gene.

Exemplary screenable markers include a  $\beta$ -glucuronidase or *uidA* gene (GUS)

which encodes an enzyme for which various chromogenic substrates are known; a  $\beta$ -lactamase gene, a luciferase gene, a xylE gene, an  $\alpha$ -amylase gene; a tyrosinase gene, an  $\alpha$ -galactosidase, or any such suitable screenable marker which is known to those of skill in the art.

5 In light of this disclosure, numerous other possible selectable and/or screenable marker genes will be apparent to those of skill in the art. Therefore, the foregoing discussion is intended to be exemplary rather than exhaustive. Although the present disclosure is exemplified in detail through the use of the *nptII* and GUS genes, the applicable techniques for making and using any other screenable or selectable marker  
10 gene will be within the skill in the art in light of the present disclosure.

The choice of the particular DNA segments to be delivered to the recipient cells will often depend on the purpose of the transformation. One of the major purposes of transformation of crop plants is to add some commercially desirable, agronomically important traits to the plant. Such traits include, but are not limited to, herbicide  
15 resistance, increased yields, insect and disease resistance, physical appearance, food content and makeup, etc. For example, one may desire to incorporate one or more genes encoding insecticidal genes which can be introduced includes the *Bacillus thuringiensis* crystal toxin gene, which provides resistance to Lepidopteran and Coleopteran insects. Such genes are well-known to those of skill in the art and are contemplated to be useful in  
20 the practice of the transformation methods disclosed herein for monocotyledonous plants such as wheat.

### 2.3 Transgenic Plants

An important aspect of the present invention are compositions comprising fertile  
25 cultivated, transgenic wheat plants. In these plants, the genome has been augmented through the genomic introduction of a preselected genetic component, to give rise to a transgenic plant. The transgene generally comprises a genetic component which includes one or more exogenous genes positioned under the control of one or more preselected genetic control elements or promoters. Such a plant is preparable by the process

described herein which includes preparing a DNA composition *in vitro* which includes the genetic component one desires to introduce into the wheat genome, then introducing the DNA into recipient wheat cells by *Agrobacterium* transformation. The genetic component typically comprises one or more of the selectable or screenable genes  
5 described above. Following transformation, wheat plants are regenerated from the cells which received the exogenous gene(s) and resulting fertile, transgenic plants may be obtained which have genomes augmented through the stable introduction of the genetic component.

In the process, the cells which are transformed may dedifferentiate, or continue to  
10 be dedifferentiated and grow in a dedifferentiated state for a period of time before undergoing differentiation and maturation into a plantlet. In preferred embodiments, the recipient cells comprise immature embryos, callus tissues, or, alternatively, suspension cells. The embryos may be freshly isolated from an immature caryopsis or, alternatively, isolated from an immature caryopsis and then treated prior to inoculation. The immature  
15 embryo may be isolated from said immature caryopsis on the same day of inoculation, or alternatively, may be one or two, or five or 10 or even more days prior to inoculation. The embryo may be an embryo that is 1 or 2 or even more days after anthesis, or alternatively, the caryopsis may be 1 or 2 or even more days post-anthesis.

When callus tissue is used as the recipient cells, the callus may be isolated from  
20 either an immature embryo or from non-embryonic cells. In certain aspects, it may be desirable to injure the embryo prior to transformation which may promote transformation efficiency.

Also disclosed is a fertile, transgenic wheat plant, whose genetic complement has been altered through the addition of a DNA composition comprising a preselected  
25 functional genetic element that includes a transgene selected from the group consisting of an *np11* gene, a *bla* gene, a *np11*, *dhfr*, *aphIV*, *aacC3*, *aacC4* gene, and a GUS gene. Particularly preferred are genes encoding glyphosate resistance, or an acceptable marker gene such as an *np11* gene.

Recipient cells may be co-transformed with more than one exogenous gene, and

under such circumstances, the exogenous genes may positioned on a single DNA segment, or alternatively, on one or more plasmids each under a different control element.

Particularly preferred plasmids are recombinant plasmids such as pMON18365, pMON32614, pMON30053, pMON25457, pMON30052 and pMON19450. The genes  
5 may be positioned under the control of a promoter such as a CaMV 35S, ubiquitin, rice actin or an enhanced CaMV 35S promoter. Where desirable, the DNA segments to be transformed may include additional 5' and/or 3' regions operatively linked to the genes.

#### 2.4 Transgenic Progeny, Seeds and Derived Cell Lines

10 Other important aspects of the invention include the progeny of the transgenic plants prepared by the disclosed methods, as well as the cells derived from such progeny, and the seeds obtained from such progeny.

### 3. BRIEF DESCRIPTION OF THE DRAWINGS

15 The drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to these drawings in combination with the detailed description of specific embodiments presented herein.

20 **FIG. 1** shows the structure of pMON18365 which is an example of the plasmid harbored in *Agrobacterium* that was used in this invention. In the T-DNA region, both GUS and *nptII* genes are driven by an enhanced 35S promoter and an intron maize HSP70 intron. The GUS gene also contains two introns.

25 **FIG. 2** shows the structure of pMON18364 which was used in the construction of pMON18365.

**FIG. 3** shows the structure of pMON18342 which was used in the construction of pMON18365.

**FIG. 4** shows the structure of pMON19476 which was used in the construction of pMON18365.

5           **FIG. 5** shows the structure of pMON18361 which was used in the construction of pMON18365.

**FIG. 6** shows the structure of pMON32614.

10           **FIG. 7** shows the structure of pMON25457.

**FIG. 8** shows the structure of pMON30053.

**FIG. 9** shows the structure of pMON30052.

15

**FIG. 10** shows the structure of pMON19450.

**FIG. 10** shows the general protocol for transformation of immature embryos using the methods disclosed herein. d represents day, w represents week.

20

**FIG. 11** shows the general protocol for transformation of suspension cells using the methods disclosed herein.

#### 4. DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

##### 25   4.1 Some Advantages of the Invention

Those having skill in the art will appreciate the many advantages of the methods and compositions provided by the present invention. A few such advantages may be summarized as follows:

#### 4.1.1 Achievement of Efficient Transformation Using Regular Binary Vector

A regular binary vector pMON18365 was used with all the experiments in this invention. Efficient transformation was achieved in most of the experiments. The fact suggested that super binary vector may not be necessary, whereas it has been shown to be essential for achieving high transformation of maize (Ishida *et al.*, 1996).

#### 4.1.2 Transformation Using Different Types of Explants

Different types of explants, immature embryos, embryogenic callus and suspension cells, were used as explants for *Agrobacterium* infection in this invention. Stable transformants (colonies from the suspension cells and plants from the embryos and embryogenic callus) were generated from all of them, which made this invention superior to other published transformation systems via *Agrobacterium*, such as the system on maize (Ishida *et al.*, 1996) in which only freshly isolated immature embryos could be used to produce transgenic plants.

#### 4.1.3 Efficient and Rapid Production of Stable Transformants

A high efficient GUS gene expression was observed among the explants shortly after the *Agrobacterium* infection as illustrated in Tables 4 and 7. In the case the suspension cells were inoculated, approximately 200 stable transformants (colonies) could be recovered from 1 ml of the cells after 40-60 d selection on a medium containing a selective agent, such as G418. Transformed plantlets could be developed from infected immature embryos and callus pieces on a G418-containing medium a few weeks after inoculation. It took only approximately 3 months from the *Agrobacterium* infection to plants in soil. Among the experiments which generated transgenic plants, the stable transformation efficiency ranged from 0.3 to 4.3% (number of transgenic events/number of explants inoculated) (Table 5), which was comparable or higher than published ones using other transformation systems (Vasil *et al.*, 1992, 1993; Weeks *et al.*, 1993; Nehra *et al.*, 1994; Becker *et al.*, 1994; Zhou *et al.*, 1995). The transgenic plants had normal morphology and completely fertile



#### 4.1.4 Mendelian Segregation of the Transgene in the Progeny

The transgene expression has been studied in the progeny ( $R_1$  generation) of 4 primary transformed plants from 2 events. The GUS activity was observed in approximately  $\frac{3}{4}$  of the individual (seeds and plants) of  $R_1$  generation (Table 6) which indicated that the gene was transmitted to the progeny in a Mendelian fashion. No abnormalities were observed among the plants.

#### 4.1.5 Production of Regenerable Transgenic Plants

A total of 9 events have been identified from 5 studies (7 different treatments) (TABLE 3). Southern analysis has been performed on 5 plants from 3 of the events from ExpAG2 and AG13. The results indicated that all 5 plants had the transgenes integrated into the genome. Plants from 6 other events (from ExpAG22, AG25 and AG30) were identified by the GUS histochemical assay and they are either in soil or in Sundae cups. All of them had very high to moderate GUS activity. The transformation efficiencies from 2 studies were 2.7 and 2.1%, respectively.

All events were generated from inoculated embryogenic callus tissue or pre-cultured immature embryos (IEs) (TABLE 4). The inoculation methods and conditions also varied among the studies. The results have indicated that *Agrobacterium*-mediated wheat transformation is repeatable and embryogenic callus tissue and pre-cultured IEs are suitable explants for *Agrobacterium* infection.

#### 4.1.6 The 3:1 Segregation Of GUS Activity In The $R_1$ Progeny

$R_1$  plants from 3 transgenic plants from ExpAG13 (TABLE 3) have been produced by germinating the  $R_1$  immature embryos *in vitro*. Plants were moved to soil, and GUS activity was in the  $R_1$  progeny, the immature seeds with the embryos removed were cut to two halves longitudinally and used for the GUS histochemical assay. Scutellum tissue was also taken for the GUS assay from some of the germinating embryos. The maternal tissue, pericarp, of each seed was GUS-positive as expected.

However, the aleurone layer of the seeds and scutellum tissue of the embryos segregated to GUS-positive or GUS-negative. The ratio of GUS-positive to GUS-negative plants was not significantly different from 3:1 by  $\chi^2$  test (TABLE 2), indicating that the transgene was most probably inserted into a single locus.

5

**TABLE 3**  
**Transgenic Events Generated by *Agrobacterium*-Mediated Transformation**

Exp-Trt	Explant	# Pieces	Inoculation (%) <sup>2</sup>	# Event (plant) <sup>1</sup>	TE
AG2-05	3-w callus	73 (large)	3 h, vacuum infiltration	2 (2)	2.7
AG13-03	2-w callus	47 (large)	vacuum inoculation	1(3)	2.1
AG22-05	10-d callus	239 (small)	3 h, soaking	1	NA
AG22-06	25-d callus	308 (small)	3 h, soaking	1	NA
AG25-24	6-d IEs	40	3 h, soaking	1	NA
AG30-02	3-d IEs	98	3 h, soaking	1	NA
AG30-05	3-d IEs	104	3 h, soaking	2	NA

TABLE 4

GUS Activity in Aleurone Layer or Scutellum in Progeny Seeds of R<sub>0</sub> Plants

R <sub>0</sub> Plant	# R <sub>1</sub> Seeds Assayed	# GUS- positive Seeds	# GUS- negative Seeds	chi square value <sup>1</sup>
AG13-03-02-01	31	22	9	0.27
AG13-03-02-02	28	22	6	0.19
AG13-03-02-03	124	96	28	0.387

<sup>1</sup> This value was calculated based on the hypothesis of 3:1 segregation. The critical value at  $\alpha=0.05$  and  $df=1$  was 3.84, larger than anyone of the chi square values in the table. Therefore, all the segregation ratios were not significantly different from hypothesized 3:1.

#### 4.1.7 Methods Comprising EPA- and FDA-Approved Selectable Markers

While many of the disclosed methods for related grains have relied on hygromycin phosphotransferase (HPT) as the selectable marker, the present invention overcomes this limitation by using *nptII* as the selectable marker.

The *nptII* gene (which encodes neomycin phosphotransferase) is the most widely-accepted selectable marker in the art, and it has been approved by the Environmental Protection Agency as a pesticidal inert. Likewise, its approval by the Food and Drug Administration as an indirect food additive makes such a marker superior to the use of markers such as hygromycin, which have not been similarly approved by these agencies.

#### 4.2 Differences from Systems Developed for Rice and Maize

The present invention differs considerably from methods described recently in other cereal grains. Unlike previous methods used for the transformation of rice, the present methods do not require high osmotic treatment for inoculation. Likewise, the present invention differs from the methods described for transformation of maize in that it

is not limited to using only freshly-isolated immature embryos as explants. The present invention also works well for pre-cultured immature embryos or callus tissue, and is not limited to freshly-isolated embryonic tissue.

Another important distinction from previous work is that the methods of the present invention are applicable using standard binary plasmids which are well-known in the art, and do not rely on the construction and use of "super" binary plasmids as disclosed in earlier reports. The binary plasmid is the most common type used for most dicotyledonous plant transformation methods, and are readily available to those of skill in the art for use in the methods disclosed herein.

\*\*\*\*\*

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

## 5. EXAMPLES

### 5.1 Example 1 — Transformation Using Immature Embryos

#### 5.1.1 Explant Preparation

A spring wheat *Triticum aestivum* cv. Bobwhite was used throughout this study. Stock plants were grown in an environmentally controlled growth chamber with 16-h photoperiod at  $800 \mu\text{mol m}^{-2}\text{s}^{-1}$  provided by high-intensity discharge (HID) Sylvania lights (GTE Products Corp., Manchester, NH). The day/night temperatures were 18/16°C. Immature caryopses were collected from the plants 14-d after anthesis. Immature embryos

(IEs) were dissected aseptically and cultured on one of the following pre-culture media before inoculation: 1) semisolid callus induction medium CM4 or CM4C (TABLE 5) for 1-6 d; 2) liquid CM4C supplemented with 0.25 M raffinose and mannitol, respectively, for 1 d; 3) liquid CM4 medium with 1/10 strength of the MS salts and supplemented with 10 g/l glucose, 3.9 g/l MES for 1 to 3 h. All the cultures were conducted at the temperature of 23-25°C.

TABLE 5  
Supplemental Components in Basic Media<sup>1</sup>

Components	CM4	CM4C	MMS.2C	MMS0C	MS1WSM <sup>4</sup>	MS2WCM <sup>4</sup>
2,4-D (mg/l)	0.5	0.5	0.2	--	1.0	2.0
Pichloram (mg/l) <sup>2</sup>	2.2	2.2	--	--	--	--
Maltose (g/l)	40	40	40	40	--	--
Sucrose (g/l)	--	--	--	--	20	20
Glutamine (g/l)	0.5	0.5	--	--	--	--
Magnesium Chloride (g/l)	0.75	0.75	--	--	0.75	0.75
Casein Hydrolysate (g/l)	0.1	0.1	--	--	--	--
MES (g/l)	--	1.95	1.95	1.95	--	--
Ascorbic Acid (mg/l) <sup>2</sup>	--	100	100	100	--	--
L-Asparagine (g/l)	--	--	--	--	0.15	0.15
Thiamine HCl (mg/l)	--	--	--	--	0.5	0.5
Gelling agent (g/l) <sup>3</sup>	2 (P)	2 (P)	2 (G)	2(G)	--	2(P)

<sup>1</sup>All media contain basal salts (MS basal salts) and vitamins (MS vitamins) from Murashige and Skoog (1962) medium unless mentioned specifically. The pH in each medium was adjusted to 5.8

<sup>2</sup>Filter-sterilized and were added to the medium after autoclaving

<sup>3</sup>Phytigel™ (P) or Gelrite® (G).

<sup>4</sup>No MS vitamins were present in the medium.

10

### 5.1.2 *Agrobacterium* Culture

Disarmed *Agrobacterium tumefaciens* strain C58 (ABI) harboring binary vector pMON18365 (FIG. 1) were used for all the studies. pMON18365 contains the Gl'S (*uidA*)

gene with an intron and *nptII* gene as a selectable marker inside the T-DNA (transfer DNA) region. Both genes were under control of an enhanced CaMV 35S promoter. Cultures of *Agrobacterium* were initiated from glycerol stocks and grown overnight at 25-26°C on a rotary shaker (150 rpm) in liquid LB medium (Miller, 1972) containing 50 mg/l each of kanamycin, streptomycin and spectinomycin, 25 mg/l chloramphenicol and 200 µM acetosyringone, to mid log phase ( $OD_{660} = 1-1.5$ ). *Agrobacterium* cells were collected by centrifugation and resuspended in an inoculation medium, CM4 or CM4C (TABLE 9) with 1/10 strength of the MS salts and supplemented with 10 g/l glucose and 200 µM acetosyringone. The *Agrobacterium* cell density was adjusted to  $OD_{660} 2$  for inoculation.

### 5.1.3 Inoculation And Co-Cultivation

The IEs temporarily maintained in the liquid pre-culture medium 3 were transferred into *Agrobacterium* cell suspension in petri dishes (25 × 100 mm). The ratio between *Agrobacterium* and IEs is about 30 ml: 200 IEs. A surfactant, Silwet® (Monsanto, St. Louis, MO), was added to the inoculation medium with a concentration of 0.01-0.075%. The inoculation was performed at 23-25°C for 3 h in dark. After inoculation, the *Agrobacterium* was removed by vacuum or using a transfer pipette, and the IEs were placed on the co-culture medium, CM4 or CM4C with 1/10 strength of the MS salts and 1.5 mg/l extra 2,4-D and supplemented with 10 g/l glucose and 200 µM acetosyringone. The embryos were placed with scutellum-side up and were co-cultured with *Agrobacterium* for 2-3 d at 25°C in the dark. The Pre-cultured immature embryos (cultured on medium 1 to 6 d) were inoculated with *Agrobacterium* with one of the following methods: 1) Immersing in the *Agrobacterium* cell suspension for 1-3 h. 2) The IEs were immersed in the *Agrobacterium* cell suspension in Petri dishes or cell culture clusters, which were then placed in a dessicator and vacuumed for 3 h using an in-house vacuum system. Silwet® at 0.01% was added to the inoculation medium in some of the studies. After inoculation, the immature embryos were blotted on sterile filter papers and then transferred to one of the following media for co-cultivation: 1) semisolid CM4 or CM4C medium (TABLE 5) supplemented with 10 g/l glucose and 200 µM of acetosyringone; 2) liquid or semisolid

CM4C medium with 1/10 strength of the MS salts and supplemented with 10 g/l glucose and 200  $\mu$ M of acetosyringone. A filter paper (Whatman No. 1) was layered on the medium in some of the studies; 3) liquid CM4 or CM4C with 1/10 strength of the MS salts and 3.9 g/l MES and supplemented with 10 g/l glucose and 200  $\mu$ M of acetosyringone. The immature embryos were placed on the medium or on the filter paper with scutellum-side up. To culture the embryos in a liquid medium, each culture plate was prepared by adding 8 ml of the liquid medium to a Petri dish (15 mm  $\times$  100 mm) containing 6 pieces of Whatman No1 filter paper (8.5 cm). The explants were co-cultured with *Agrobacterium* at 24°C in the dark for 2 to 3 d. After co-culture, the immature embryos were rinsed with liquid CM4 or CM4C medium supplemented with 500 mg/l carbenicillin (delay medium). The infected immature embryos were cultured on the solid delay medium for 2-5 d. The general protocol for these transformation methods is given in FIG. 6.

#### 5.1.4 Efficiency of T-DNA Delivery

The efficiency of T-DNA delivery was measured by transient GUS expression assay after 2-3 d delay of selection. High levels of transient GUS expression were observed for most of studies, indicating that the T-DNA delivery was very efficient. The effect of Silwet® on the T-DNA delivery when included in the inoculation medium was extensively investigated on various explants. As shown in TABLE 6, Silwet® at 0.05-0.1% significantly enhanced the transient GUS expression on the freshly isolated IEs. Similar results were also observed using pre-cultured IEs and embryogenic callus as explant for inoculation. For the immature embryos pre-cultured for 1 or 2 d, the GUS spots were located mostly along the edge of the scutellum from which less embryogenic callus developed from. The GUS spots, however, were found in the callusing areas on the embryos pre-cultured 3 d or longer.



TABLE 6

Effect of Surfactant on GUS Expression When Present in Inoculation Medium<sup>1</sup>

Silwet (%)	GUS-positive spots/explant	No. of explants w/GUS-positive spots/total explant (%)
0.00	7.8	11/34 (32)
0.01	17.6	15/19 (79)
0.05	149	13/13 (100)
0.1	111	8/8 (100)
0.5	1	1/1 (100)

<sup>1</sup> The explants were IEs isolated a few of hours before inoculation.

5

#### 5.1.5 Selection And Plant Regeneration

After 2 to 5 d on the delay medium, the *Agrobacterium*-infected immature embryos were transferred to the callus induction medium, CM4 or CM4C (TABLE 5) medium with 25 mg/l G418 and 250 mg/l carbenicillin. The immature embryos were cultured for 2 to 3 weeks for callus induction before being transferred to the first regeneration medium, MMS.2C (TABLE 5) with 25 mg/l G418 and 250 mg/l carbenicillin. At transfer to the regeneration medium, each piece of callus was divided into several small pieces (~2 mm). Two weeks after the cultures on the first regeneration medium, young shoots and live callus tissue were transferred to the second regeneration medium, MMS0C (TABLE 5) with the same concentrations of G418 and carbenicillin. Plantlets, which were confirmed later true transformants, grew vigorously and formed strong root systems in this medium. However, there were some plantlets showing some resistance to G418 at this stage. They could grow well and form one or a few of roots, although they did not grow as vigorously as the true transformants. When the plants were about 3 cm or longer, they were transferred to Sundae cups (Sweetheart Cup Company, Chicago, IL) containing the second regeneration medium as above for further growth and selection. Leaf samples were taken from some of the

10

15

20

plantlets for the GUS histochemical assay at this stage. However, the plantlets, which appeared resistant and showed no GUS activity at this stage, were not eliminated. During growth in the cups, most of the non-transformants died or showed signs of susceptibility to G418. The plants highly resistant to G418 (growing vigorously with strong root system) were moved to soil before they reached the top of the cups. All the plants originated from the same embryo were considered as siblings from the same event.

#### 5.1.6 Confirmation Of The Transgenic Nature Of The Plants

The plants were grown in a environmentally controlled growth chamber under the same growth conditions as described above. Since it took only approximately 3 months from inoculation to transferring most of the plants to soil, no visible abnormalities, which are usually associated with plants having prolonged *in vitro* culture period, were observed among the plants. The plants were completely fertile. Each plant was examined by one or more of the following methods:

- (1) The GUS histochemical assay (Jefferson, 1987) using different parts of the plants.
- (2) Biological assay (leaf bleach assay). Before heading, leaf samples (~5-7 mm long) were taken from the youngest fully expanded leaves and were placed in wells of 24-well cell culture clusters (Costar Corporation, Cambridge, MA). Each well was filled with 0.5 ml water solution composed of 300 mg/l paromomycin (Sigma) and 100 mg/l Benlate (a fungicide made by Du Pont) or 100 mg/l Benlate alone. Three leaf samples from the same leaf of each plants were placed in 2 wells containing paromomycin and Benlate and 1 well containing Benlate alone, respectively. Leaf samples from the non-transformed Bobwhite plants were used as negative controls. The samples were vacuum-infiltrated in a dessicator using an in-house vacuum system for 5 min and then the cluster were sealed very well with Parafilm<sup>®</sup> before being placed under light ( $140 \mu\text{mol m}^{-2}\text{s}^{-1}$ ). The results were determined 60 h later. The leaf samples which were highly resistant to paromomycin remained green in most area except the two edges (~1 mm wide), which indicated that the plants had the functional *hptII* gene. The leaf samples from the plants without the gene or

with the non-functional gene were bleached out completely by paromomycin as the negative controls, or had only small patches of green areas.

(3) Southern hybridization analysis (Southern, 1975). Genomic DNA were isolated from leaf tissue of the plants following the method of Shure *et al.* (1983). Fifteen  
5  $\mu\text{g}$  of genomic DNA was digested with restriction endonuclease *Bam*HI and fractionated on a 0.8% agarose gel. The DNA was transferred to Hybond N membranes (Amersham, Arlington Heights, IL) according to standard procedures (Sambrook *et al.*, 1989). The probe for detecting the *nptII* gene was prepared by gel purifying a 977 bp *Nco*I fragment from plasmid pMON18365 (FIG. 1). The fragment was labeled with  $^{32}\text{P}$  dCTP using  
10 random primer labeling kit (Prime-It II<sup>®</sup> from Stratagene<sup>®</sup>, La Jolla, CA), to a specific activity of  $2.6 \times 10^9$  cpm/ $\mu\text{g}$ . The membrane was hybridized for 14 h at 42°C in a solution containing 50% formamide, 5× SSC, 5× Denhardt's, 0.5% SDS, 100  $\mu\text{g}/\text{ml}$  tRNA. The condition of the final wash was 0.1% SSC and 0.1% SDS at 60°C for 15 min.

#### 15 5.1.7 Efficiency Of Stable Transformation

The number of the transgenic events in each study was determined after the plants were assayed as described above. The transformation efficiency (number of events/ number of immature embryos) varied from study to study and among different treatment conditions that generated transgenic plants (TABLE 7). However, they were comparable with or  
20 higher than any published wheat transformation efficiencies (Vasil *et al.*, 1992, 1993; Weeks *et al.*, 1993; Nehra *et al.*, 1994; Becker *et al.*, 1994; Zhou *et al.*, 1995).

TABLE 7

## Stable Transformation Efficiencies in the Production of Transgenic Wheat

Experiment- Treatment	Explant	Number of IEs or callus pieces (A)	Number of transgenic events (B) <sup>1</sup>	Efficiency (B/A%)
AG2-05	21-d callus, intact	73	2	2.7
AG13	14-d callus, intact	47	1	2.1
AG22-05	10-d callus, ~2 mm	239	1	0.4
AG22-06	25-d callus, ~2 mm	308	1	0.3
AG22-11	10-d callus, ~2 mm	232	1	0.4
AG25-24	6-d IEs	40	1	2.5
AG27-15	1-d IEs	23	1	4.3
AG29-04	5-d IEs	97	1	1.0
AG30-02	3-d IEs	98	1	1.0
AG30-05	3-d IEs	104	2	1.9
AG30-08	3-d IEs	36	1	2.8
9528	0-d IEs	160	1	0.6
9531	17-d callus, intact	50	1	2.0
9602	0-d IEs	250	3	1.2
9604	0-d IEs	700	1	0.14
9608	0-d IEs	124	1	0.8
9609	0-d IEs	140	2	1.4
9614	0-d IEs	38	1	2.6
9620	15-d callus, intact	110	3	2.7

<sup>1</sup> Each transgenic event had one or more plants.

### 5.1.8 Progeny Analysis Of The Transgenic Plants

Immature caryopses were harvested from the primary transgenic plants ( $R_0$  generation) with both *nrp1II* and GUS activities approximately 20 days after anthesis. The immature embryos were isolated and cultured on MMSOC medium (TABLE 5) for plant germination (approximately 1 week in Petri dishes and another week in Sundae cups with the same medium). Each of the immature caryopses with the embryos removed was cut to two halves longitudinally and moved to a well in 96-well cell culture clusters (Costar Corp., Cambridge, MA) for the GUS histochemical assay as described above. All the half caryopses were examined under microscope to determine the GUS activity in different areas. Part of the scutellum tissue was also taken from each germinating embryo for the GUS assay. After the plants ( $R_1$  generation) were moved to soil approximately 2 weeks after germination, the GUS activities in the plants were determined by the GUS histochemical assay using leaf and flower tissues, respectively.

As expected, the maternal tissue, pericarp, of each immature caryopses showed GUS activity. However, in the aleurone layer most of the caryopses showed GUS activity, others not. The ratio of caryopses with GUS positive aleurone layer to ones with GUS negative aleurone layer was not significantly different from 3:1 by chi square test (TABLE 8). The data from the GUS assay on the caryopses matched well with the data on scutellum tissue and later with the GUS assay on leaf and flower tissue, although the GUS activity in aleurone layer and scutellum appeared much stronger than in leaf and flower tissues.

**TABLE 8**  
**Segregation of GUS Activity in R<sub>1</sub> Seeds and Plants**

R <sub>0</sub> plants	Number of R <sub>1</sub> seeds (plants) assayed	Number of GUS-positive	Number of GUS-negative	chi square value <sup>1</sup>
16612	31	22	9	0.27
16613	28	22	6	0.19
16614	124	96	28	0.387
16953	128	103	25	2.04

<sup>1</sup> This value was calculated based on the hypothesis of 3:1 segregation. The critical value at  $\alpha=0.05$  and  $df=1$  was 3.84, larger than any one of the chi square values in the table. Therefore, all the segregation ratios were not significantly different from hypothesized 3:1.

## 5.2 Example 2 -- Transformation Using Embryogenic Callus

### 10 5.2.1 Explant Preparation

Immature embryos of wheat (*Triticum aestivum* L.) cv Bobwhite were isolated from the immature caryopsis 14 days after anthesis and cultured on callus induction medium CM4 or CM4C (TABLE 9) with scutellum-side up. After 10 days or longer, immature embryos developed into embryogenic callus. Each callus piece was approximately 5 mm or bigger. The callus pieces were inoculated with *Agrobacterium* without being broken down (intact), or only the most embryogenic callus sections were selected and broken down into small pieces (~2 mm) using fine-pointed forceps for inoculation.

### 5.2.2 Inoculation And Co-Cultivation

20 Callus pieces were inoculated with the *Agrobacterium* cell suspension prepared as described above, using one of the following methods:

- 1) Immersing the callus pieces in the *Agrobacterium* cell suspension for 3 h.
- 2) Immersing the callus pieces in the *Agrobacterium* cell suspension with

vacuum infiltration for 3 h.

3) Intact callus pieces were placed on a piece of sterile filter paper saturated with the liquid inoculation medium. The callus pieces were inoculated as they were, or were pressed before inoculation using a sterile spatula until the callus pieces became "pancakes". Each piece of the filter paper holding the callus pieces was transferred to a 150-ml filter system (Corning, Inc., Corning, NY) connected to an in-house vacuum system, or to a Buchner funnel connected to the vacuum system through a filter flask. With the vacuum system on, the *Agrobacterium* cell suspension was dropped onto the callus pieces (>1 ml over 6 callus pieces) slowly. After dropping the *Agrobacterium* suspension cells, vacuum was applied for another 10 min. The callus pieces inoculated with any one of the methods were transferred to Petri dishes (100 × 15 mm) each containing 6 pieces of Whatman filter paper (No. 1, 8.5 cm) saturated with 8 ml of the co-cultivation medium (CM4 or CM4C with 1/10 strength of the MS salts and supplemented with 10 g/l glucose and 200 µM acetosyringone). The plates were sealed well with Parafilm<sup>®</sup>. After co-culture with the *Agrobacterium* cells for 3 days, the callus pieces were washed with the liquid delay medium (CM4 or CM4C supplemented with 500 mg/l carbenicillin) and blotted on pieces of sterile filter paper to remove excess liquid. The callus pieces were cultured on the semisolid delay medium for 3 days before being transferred to the selective callus induction medium.

20

### 5.2.3 Efficiency Of T-DNA Delivery

After co-culture with *Agrobacterium* or delay stage, samples of the callus pieces were randomly taken for the GUS histochemical assay (Jefferson, 1987). As shown in TABLE 11, most callus pieces assayed had GUS positive spots. In some studies (such as experiment AG13), every piece of callus assayed had GUS positive spots. The number of GUS positive spots in each piece of callus varied from a few of to approximately 100. In experiment AG13, another GUS histochemical assay was carried out two weeks after inoculation. All of the 16 pieces of callus assayed had some GUS positive spots and among them 7 pieces (44%) had growing GUS positive sectors, which suggested that transformed

25

cells were proliferating.

TABLE 9

Efficiency of T-DNA Delivery as Indicated by GUS Histochemical Assay on Callus

5

Pieces After Co-Culture

Exp- Treatment	Explant	Number of IEs w/ GUS-positive spots (Total number of IEs assayed)	Number of spots on each GUS-positive IE
AG2-5	21-d callus, intact	1/19	1 - 5
AG13	14-d callus, intact	25/25	several - ~50
AG22-5	10-d callus, ~2 mm	6/15	several
AG22-6	25-d callus, ~2 mm	7/10	a few - ~100
AG22-11	10-d callus, ~2 mm	12/20	a few - dozen

#### 5.2.4 Selection And Plant Regeneration

The callus pieces, which were intact during inoculation, were cultured on the selective callus induction medium (CM4C supplemented with 25 mg/l G418 and 250 mg/l carbenicillin) as they were, or were broken down to several small pieces (~2 mm) at transfer to the medium. The inoculated small callus pieces were left intact and cultured on the medium. After 2 to 3 weeks, callus pieces were transferred to the regeneration medium. The regeneration procedure was the same as described in Example 1.

#### 15 5.2.5 Transformation Efficiency

The regenerated plants showed no visible abnormalities and were completely fertile. All the plants were tested with the biological assay, GUS histochemical assay and/or Southern hybridization assay as described in Example 1. A total of 10 transgenic events have been generated from 7 experimental treatments using embryogenic callus at different



ages (TABLE 9). The transformation efficiency (number of events/number of the callus pieces) ranged from 0.3 to 2.7%.

### 5.2.6 Progeny Analysis Of The Transgenic Plants

5 The R1 plants were analyzed as described in Example 1.

## 5.3 Example 3 -- Transformation of Suspension Cells

### 5.3.1 Explant Preparation

10 Wheat v. Mustang suspension cells were cultured in liquid MSIWSM (TABLE 9) at 28°C in the dark on a rotary shaker (250 rpm). Cells harvested after 3 d subculture were used for inoculation with *Agrobacterium*. The general protocol for transformation of suspension cells is given in FIG. 7.

### 5.3.2 *Agrobacterium* Culture and T-DNA Delivery

15 The *Agrobacterium* culture method was essentially the same as described in Example 1. The *Agrobacterium* cell density was adjusted to OD<sub>660</sub> 0.5-1 in inoculation medium (CM4 medium with 1/10 strength of the MS salts and 3.9 g/l MES and supplemented with 10 g/l glucose and 200 µM acetosyringone. Liquid medium was removed from the wheat suspension culture by vacuum. Each ml of the wheat cells was  
20 mixed with 3 ml of the *Agrobacterium* cell suspension petri dishes (100 × 25 mm). The inoculation was performed at 23-25°C for 30 min to 4 h. After inoculation, *Agrobacterium*-infected wheat cells were placed on a piece of sterile Whatman filter paper in petri dishes (100 × 15 mm). The filter paper was wetted with the liquid medium as described above. The co-culture plates were placed in the dark at 23-25°C for 2 to 3 d.

25

### 5.3.3 Recovery of Transformed Colonies

After co-culture, the inoculated cells were transferred into liquid MS2WCM (TABLE 9) with 500 mg/l carbenicillin and cultured in flasks for 1 d with gentle agitation, and then plated on a piece of filter paper on the solid MS2WCM medium supplemented

with either 25 mg/l G418 or 50 mg/l paromomycin and 250 mg/l carbenicillin for selection. Resistant colonies could be recovered after 40 to 60 d selection. The transformation was highly efficient based on the number of transformed colonies recovered. Routinely approximately 200 independently transformed colonies were recovered from 1 ml of inoculated cells.

#### 5.3.4 Factors Influencing Transformation Efficiency

Three factors involved in inoculation and co-culture processes were found to significantly influence the transformation efficiency of suspension cells. The factors were inoculation and co-culture temperatures, inoculation and co-culture time period, and *Agrobacterium* cell density for inoculation. As shown in TABLE 10, the best inoculation and co-culture temperature is 23 to 25°C, and the transformation efficiency is significantly reduced at temperature 19°C or 28°C. The best inoculation time period was 30 min (TABLE 11), and high transformation efficiency could be achieved with 2 or 3 d co-culture. The co-culture time period of 1 d or shorter significantly reduced the transformation efficiency. The best *Agrobacterium* cell density for inoculation was OD<sub>660</sub> 0.5 (TABLE 12). *Agrobacterium* density higher or lower than that reduced the transformation efficiency.

TABLE 10  
Effect of Temperature During Inoculation and Co-Cultivation on Stable Transformation of Suspension Cells

Temperatures (°C)	Number of GUS-positive colonies/ml cell inoculated
19	52.5 ± 19.36
23	290 ± 70.83
25	273 ± 65.43
28	132.5 ± 22.6

TABLE 11

Effect of Inoculation and Co-Culture Time on Suspension Cell Transformation

Stage	Duration (h)	Number of GUS-positive colonies/ml cells inoculated
Inoculation	0.5	351.25 $\pm$ 29.23
	1	178.5 $\pm$ 23.1
	2	186.75 $\pm$ 23.10
	3	202 $\pm$ 51.37
	4	152.25 $\pm$ 29.68
	14	4 $\pm$ 0.82
Co-culture	24	53 $\pm$ 4.24
	48	232.5 $\pm$ 26.29
	72	258.5 $\pm$ 15.26

5

TABLE 12

Effect of *Agrobacterium* Cell Density on Transformation of Suspension Cells

<i>Agrobacterium</i> cell density (OD <sub>660</sub> )	Number of GUS-positive colonies/ml of cells
0.10	79 $\pm$ 4.97
0.25	174 $\pm$ 53.67
0.50	260.25 $\pm$ 45.33
1.00	172 $\pm$ 51.04
2.00	159 $\pm$ 70.06

### 5.3.5 Confirmation of Transformed Colonies

The transformation was confirmed by histochemical GUS assay as described. Samples from 51 G418 resistant colonies were assayed for GUS activity. Forty nine out of 51 colonies showed GUS positive. Among them, fifteen were sampled for Southern blot analysis. All the samples showed a strong hybridization signal to the *nptII* probe. These results may suggest that the co-expression frequency was close to 100%.

## 5.4 Example 4 – Transformation of Suspension Cells

### 5.4.1 *Agrobacterium* Constructs

Plant transformation vectors similar to those derived from a Ti plasmid of *Agrobacterium tumefaciens*, such as those disclosed by Herrera-Estrella *et al.* (1983), Bevan *et al.* (1983), Klee *et al.* (1985) and EPO publication 120,516 (Schilperoort *et al.*) were constructed for use in wheat. The *Agrobacterium* binary vector for wheat, pMON18364 (FIG. 2), was constructed by the removal of the P-nos/*nptII*/nos 3' from pMON18342 (FIG. 3), by digesting pMON18342 with *NotI* and *HindIII* restriction enzymes to completion and isolation of the 5.7 kb fragment. This resulting 5.7 kb vector backbone fragment contains the ori-322 replication origin for replication in *E. coli* and the ori-V region for replication in *Agrobacterium*, bacterial resistance markers for spectinomycin and streptomycin selection. The P-e35S/hsp70 intron/*kan*/nos 3' chimeric fragment was isolated from pMON19476 (FIG. 4) by digesting to completion with *NotI* and *HindIII* restriction enzymes and gel isolation of the resulting 2.7 kilobase (kb) insert. pMON18364 the binary vector was constructed by ligation of the 2.7 kb *NotI*, *HindIII* P-e35S/hsp70 intron/*kan*/nos 3' fragment and the *NotI*, *HindIII* 5.7 kb vector backbone fragment using T4 DNA ligase. Ligation products were transformed into Novablu cells and selected for spectinomycin resistance. Transformed colonies from this unidirectional cloning were grown in liquid culture and DNA was analyzed for the presence of the P-e35S/*kan*/nos 3' chimeric gene by restriction digestion with *HindIII*, and *EcoRV*. For reporter gene constructs, the *Agrobacterium* binary vector containing the  $\beta$ -glucuronidase (GUS) gene was constructed by digesting pMON18361 (FIG. 5) with *HindIII* and isolation

of the 3.8 kb fragment containing the P-e35S/hsp70 intron/GUS/nos 3' chimeric fragment and ligation of this fragment to *Hind*III digested pMON18364. Analysis of the resulting transformed colonies was performed with *Hind*III and orientation was verified with *Bgl*II, and *Xba*I digestions. The resulting construct is pMON18365 (FIG. 6): RE>P-e35S/hsp70  
5 intron/GUS:ST-LS1 intron/nos 3'; P-e35S/HSP70 intron/nptII/nos >LB, where LB and RB are left and right *Agrobacterium* Ti plasmid transfer borders, respectively.

Other genes may be introduced in wheat; for example, wheat streak mosaic virus (WSMV) coat protein and Barley Yellow Dwarf Virus (BYVD) replicase conferring resistance to plant viruses. A suitable plant expression vector is pMON18365 or  
10 derivatives of similar binary vectors. A gene encoding the coat protein (CP) from WSMV may similarly be inserted into pMON18364 or pMON18365 using standard techniques. The recombinant vector containing the WSMV CP gene when transferred into wheat plants by the *Agrobacterium* mediated transformation method would be expected to confer resistance to the wheat plants from infection by WSMV. Similarly, a  
15 gene encoding the BYVD full-length replicase gene may be introduced into the same or similar *Agrobacterium* vector cassettes. The recombinant vector containing the BYDV full-length replicase gene when transferred into wheat plants by an *Agrobacterium* mediated transformation method would be expected to confer resistance to the wheat plants from infection by BYDV.

20 Genes encoding fungal resistance may also be introduced into *Agrobacterium* binary vectors such as pMON18364 and pMON32614 (FIG. 6). Antifungal protein genes such as those encoding Alfalfa or *Alyssum* antifungal proteins inserted into pMON18364 or pMON18365 as described above would allow engineering for fungal resistance into wheat by *Agrobacterium* mediated transformation for the production of transgenic plants.  
25 The resulting plants would be expected to exhibit resistance to *Fusarium* or other fungal or bacterial pathogens.

Genes that affect quality traits involved in carbohydrate quantity and composition in the kernel such as ADP glucose pyrophosphorylase (ADPGPP) expressed under the control of a kernel enhanced promoter or other tissue enhanced promoters could be

inserted into a binary vector such as pMON18365 or derivatives  
*Agrobacterium*-mediated transformation of wheat plants with this gene are expected to  
confer changes in the amount or quality of carbohydrates in the kernel or other tissues  
producing changes in the composition of matter.

5           One may also insert genes affecting traits in wheat such as tolerance to herbicides  
for herbicide resistance, tolerance to drought and high salinity, and tolerance to cold and  
heat stresses. Genes increasing yield of wheat plants, genes used for male sterility for the  
production of hybrid wheat, and genes influencing germination could also be introduced  
into binary plant transformation vectors for *Agrobacterium*-mediated transformation  
10       producing wheat plants expressing traits of the desired phenotype. Examples of plasmids  
useful for inserting genes include pMON25457 (FIG. 7), pMON30053 (FIG. 8),  
pMON30052 (FIG. 9), and pMON19450 (FIG. 10).

## 15       6.       REFERENCES

The references listed below and all references cited herein are incorporated herein  
by reference to the extent that they supplement, explain, provide a background for, or  
teach methodology, techniques, and/or compositions employed herein.

- 20       U. S. Patent No. 5,179,022.  
      Intl. Pat. Appl. Publ. No. WO 91/02071.  
      Intl. Pat. Appl. Publ. No. WO 92/06205.  
25       Intl. Pat. Appl. Publ. No. WO 93/18168.  
      Intl. Pat. Appl. Publ. No. WO 94/0077.  
30       Intl. Pat. Appl. Publ. No. WO 94/00583.  
      Intl. Pat. Appl. Publ. No. WO 95/06722.  
      Barcelo *et al* , *Plant J* , 5:583-592, 1994.

- Becker *et al.*, *Plant J.*, 5:299-307, 1994.
- Bevan *et al.*, *Nature*, 303:209, 1983
- 5 Chan *et al.*, *Plant Cell Physiol.*, 33(5):577-583, 1992.
- Chan *et al.*, *Plant Mol. Biol.*, 22(3):491-506, 1993.
- 10 Chan *et al.*, *Plant Physiol.*, (Rockville) 108 (2 Suppl.) 115, Abstr. Annu. Meet. Amer. Soc. Plant Physiol., Charlotte, NC, 1995.
- Chen and Dale, *Transgenic Res.*, 1(2):93-100, 1992.
- 15 Chilton, *Proc. Natl. Acad. Sci. USA*, 90(8):3119-3120, 1993.
- Christou *et al.*, *Plant Physiol.*, 87:671-674, 1988.
- Christou *et al.*, *Bio/Technology*, 9:957-962, 1991.
- 20 Conger *et al.*, *Plant Cell Rep.*, 6:345-347, 1987.
- Conner and Dommissie, *Int. J. Plant Sci.*, 153:550-555, 1992.
- 25 Creissen *et al.*, *Plant Cell Rep.*, 8(11):680-683, 1990.
- Datta *et al.*, *Bio/Technology*, 8:736-740, 1990.
- Davey *et al.*, *J. Exp. Bot.*, 42:1129-1169, 1991.
- 30 De la Pena *et al.*, *Nature*, 325:274-276, 1987.
- Dekeyser *et al.*, *Plant Physiol.*, 90:217-223, 1989.
- 35 Delbreil *et al.*, *Plant Cell Rep.*, 12(3):129-132, 1993.
- Della-Cioppa *et al.*, *Bio/Technology*, 5:579-584, 1987.
- Deng *et al.*, *Sci. China Ser B Chem. Life Sci. Earth Sci.*, 33(1):27-34, 1990.
- 40 Du *et al.*, *Genet. Manip. Plants*, 5:8-12, 1989.
- Fromm *et al.*, *Proc. Natl. Acad. Sci. USA*, 82:5824-5828, 1985.

- Fromm *et al.*, *Nature*, 319:791-793, 1986.
- Fromm *et al.*, *Bio/Technology*, 8:833-839, 1990.
- 5 Godwin and Chikwamba, xiii-174, Plenum Press: New York, NY.
- Gordon-Kamm *et al.*, *Plant Cell*, 2:603-618, 1990.
- 10 Gould *et al.*, *Plant Physiol.*, 95(2):426-434, 1991.
- Graves and Goldman, *Plant Mol Biol.*, 7(1):43-50, 1986.
- Grimsley *et al.*, *Nature*, 325:177-179, 1987.
- 15 Hayakawa *et al.*, *Proc. Natl Acad. Sci. USA*, 89:9865-9869, 1992.
- Herrera-Estrella *et al.*, *Nature*, 303:209, 1983.
- 20 Hess *et al.*, *Plant Sci.*, 72(2):233-244, 1990.
- Hiei, Y., *Plant J.*, 6(2):271-282, 1994.
- Hood *et al.*, *Plant Physiol.*, 105(1 Suppl.), 114, 1994.
- 25 Ishida *et al.*, *Plant Physiol.*, (Rockville) 108 (2 Suppl.) 114, Abstr. Annu. Meet. Amer. Soc. Plant Physiol., Charlotte, NC, 1995.
- Ishida, Y., *et al.*, *Nature Biotech.*, 745-750, 1996.
- 30 Jaehne *et al.*, *Euphytica*, 85:1-3:35-44, 1995.
- Jefferson, *Plant Mol. Biol. Rep.*, 5:387-405, 1987.
- 35 Jefferson *et al.*, *EMBO J.*, 6:3901-3907, 1987.
- Kasha *et al.*, *Gene Manip. Plant Improv. II*, 213-239, 1990.
- Klee *et al.*, *Bio/Technology*, 3:637, 1985.
- 40 Knutson *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:2624-2628, 1992.
- Kornienko *et al.*, *Pushchino*, 20-22, noyabr'a, Tezisy dokladov, 20:21, 128-129, 1991.



- Langridge *et al.*, *Plant J.*, 2(4):631-638, 1992.
- Li *et al.*, *Sci. Shina Ser B Chem. Life Sci. Earth Sci.*, 34(1):54-63, 1991.
- 5 Liu *et al.*, *Plant Mol. Biol.*, 20(6):1071-1087, 1992.
- Luo and Wu, *Plant Mol. Biol. Rep.*, 6:165-174, 1988.
- 10 Mahalakshmi and Khurana, *J. Plant Biochem. Biotech.*, 4(2):55-59, 1995.
- May *et al.*, *Bio/Technology*, 13:486-492, 1995.
- McElroy *et al.*, *Mol. Gen. Genet.* 231:150-160, 1991.
- 15 Miljus-Djukic *et al.*, *Plant Cell Tissue Organ Culture*, 29(2):106-108, 1992.
- Miller, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1972.
- 20 Mooney and Doodwin, *Plant Cell Tissue Organ Culture*, 25:209-218, 1991.
- Mursahige and Skoog, *Physiol. Plant*, 15:473-497, 1962.
- Nehra *et al.*, *Plant J.*, 5:285-297, 1994.
- 25 Paszkowski *et al.*, *EMBO J.*, 3:2717-2722, 1984.
- Paszkowski *et al.*, *Plant Mol. Biol.*, 6:303-312, 1986.
- 30 Picard *et al.*, VIIth International Wheat Genetics Symposium, University of Cambridge, 779:781, 1988.
- Piorer *et al.*, *Science*, 256:520-523, 1992.
- 35 Potrykus, I., *Bio/Technology*, 8:535-543, 1990.
- Raineri *et al.*, *Bio/Technology*, 8:33, 1990.
- Raineri *et al.*, *Proc. Natl. Acad. Sci. USA*, 90(8):3549-3553, 1993.
- 40 "Report on the Use of Antibiotic Resistance Markers in Genetically Modified Food Organisms," Advisory Committee on Novel Foods and Processes, July 1994.

- Rhodes *et al.*, *Science*, 240:204-207, 1988.
- Ritchie *et al.*, *Transgenic Res.*, 2(5):252-265, 1993.
- 5 Sambrook *et al.*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Sanford, *Trends Biotechnology*, 6:299-302, 1988.
- Sanford, J.C., *Physiologia Plantarum*, 79:206-209, 1990
- 10 Sautter *et al.*, *Bio/Technology*, 9:1080-1085, 1991.
- Schlappi and Hohn, *Plant Cell*, 4(1):7-16, 1992
- 15 Schledzewski and Mendel, *Transgenic Research*, 3:249-255, 1992
- Schrott, M., *In Gene Transfer in Plants*, Springer, Verlag, New York, 1995, pp. 325-331
- Selichi *et al.*, *Plant Physiol.*, 100:1503-1507, 1992
- 20 Shen and Hohn, *Plant J.*, 2(1):35-42, 1992.
- Shen *et al.*, *Proc. Natl. Acad. Sci. USA*, 90(4):1488-1492, 1993.
- 25 Shen *et al.*, *Meth. Mol. Biol.*, 44:Xiv-417, 1995.
- Shillito *et al.*, *Bio/Technology*, 3:1099-1103, 1985.
- Shimamoto *et al.*, *Nature*, 338:274-276, 1989.
- 30 Shure *et al.*, *Cell*, 35:225-233, 1983.
- Smith *et al.*, *In Vitro Cell Develop. Biol.*, 30A (3 Part 2), 1994.
- 35 Smith and Hood, *Crop Sci.*, 35:301-309, 1995.
- Somers *et al.*, *Bio/Technology*, 10:1589-1594, 1992.
- Southern, *J. Mol. Biol.*, 98:503-517, 1975.
- 40 Topfer *et al.*, *Plant Cell*, 1:133-139, 1989.
- Vasil *et al.*, *Bio/Technology*, 10:667-674, 1992

- Vasil *et al.*, *Bio/Technology*, 11:1153-1158, 1993.
- Vijayachandra *et al.*, *Plant Mol. Biol.*, 29(1):125-133, 1995.
- 5 Wan and Lemaux, *Plant Physiol.*, 104:37-48, 1994.
- Weeks *et al.*, *Plant Physiol.*, 102:1077-1084, 1993.
- 10 Wilmink *et al.*, *Plant Cell Rep.*, 11:76-80, 1992.
- Xu *et al.*, *Chin J. Bot.*, 2(2):81-87, 1990.
- Zaghmout and Trolinder, *Nucl. Acids Res.*, 21(4):1048, 1993.
- 15 Zaghmout, *Theoret. Appl. Genet.*, 89(5):577-582, 1994.
- Zhou and Konzak, *Crop Sci.*, 29:817-821, 1989.
- 20 Zhou *et al.*, *Plant Cell Tissue Organ Culture*, 30:78-83, 1992.
- Zhou *et al.*, *Plant Cell Rep.*, 15:159-163, 1995.
- 25 Ziauddin *et al.*, *Plant Cell Rep.*, 11:489-493, 1992.

All of the compositions and methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred  
30 embodiments, it will be apparent to those of skill in the art that variations may be applied to the composition, methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same  
35 or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

## WHAT IS CLAIMED IS:

1. A fertile, transgenic wheat plant, the genome of which has been altered through the genomic introduction of a pre-selected genetic component, said component  
5 comprising an exogenous gene positioned under the control of one or more pre-selected genetic control elements, the plant prepared by a process comprising:

10 (a) preparing a DNA composition *in vitro*, which composition includes the genetic component one desires to introduce into the genome of a wheat plant;

(b) introducing said DNA composition into recipient wheat cells by *Agrobacterium* transformation;

15 (c) regenerating wheat plants from said cells which have received said genetic component; and

20 (d) identifying a fertile, transgenic wheat plant whose genome has been altered through the stable introduction of said genetic component.

2. The fertile, transgenic wheat plant of claim 1 wherein the genetic component of step (a) comprises a selectable or screenable gene.

25 3. The fertile, transgenic wheat plant of claim 1, wherein said recipient cells comprise an immature embryo, a callus tissue, or suspension cells.

4. The fertile, transgenic wheat plant of claim 3, wherein said embryo is isolated from an immature caryopsis or isolated from an immature caryopsis and then precultured prior to inoculation.

5

5. The fertile, transgenic wheat plant of claim 4, wherein said immature embryo is isolated from said immature caryopsis within from about 24 hours to about ten or more days prior to inoculation.

10

6. The fertile, transgenic wheat plant of claim 3, wherein said callus tissue is isolated from an immature embryo or developed from embryogenic cells.

15

7. The fertile, transgenic wheat plant of claim 3, wherein said immature embryo comprises an embryo 2 or more days after anthesis or said immature caryopsis comprises a caryopsis 2 or more days after anthesis.

20

8. A fertile, transgenic wheat plant, the genetic complement of which has been altered through the addition of a DNA composition comprising a pre-selected functional genetic element that includes a transgene selected from the group consisting of an *nptII* gene, a *bla* gene, a *nptI* gene, a *dhfr* gene, a *aphIV* gene, a *aacC3* gene, a *aacC4* gene and a GUS gene, wherein said functional genetic element confers on said wheat plant a phenotypic trait that is not found in the parentage of said plant.

25

9. The fertile, transgenic wheat plant of claim 1 or claim 8 wherein said gene comprises an *nptII* gene.

10. The transgenic wheat plant of claim 1 or claim 8, wherein at least two exogenous  
genes are positioned on the same DNA segment, and said recipient cells are transformed  
5 with said segment.

11. Progeny of the plant of claim 1 or 8.

10

12. Cells obtained from the plant of claim 1 or 8.

13. Seeds obtained from the plant of claim 11

15

14. A method for producing a fertile transgenic wheat plant, comprising the steps of:

- 20 (a) establishing a regenerable culture from a wheat plant to be transformed;
- (b) introducing a DNA composition comprising a genetic component one  
desires to introduce into the genome of said wheat plant, by  
*Agrobacterium* transformation;
- 25 (c) identifying or selecting a transformed cell line; and
- (d) regenerating a fertile transgenic wheat plant therefrom, wherein said DNA  
is transmitted through a complete sexual cycle of said transgenic plant to  
its progeny, wherein said progeny comprises a selectable or screenable

marker gene, and wherein said marker gene is chromosomally integrated.

15. The method of claim 14, wherein at least two exogenous genes are positioned on  
5 the same DNA segment, and said regenerable culture is transformed with said segment.

16. The method of claim 14, wherein said *Agrobacterium* is *A. tumefaciens* C58.

10

17. The method of claim 14, wherein said culture comprises an immature embryo, a  
callus tissue, or suspension cells.

15 18. A method for producing a transgenic wheat plant, comprising the steps of

(a) establishing a culture from a wheat plant to be transformed;

20 (b) transforming said culture with an *Agrobacterium* comprising a DNA  
composition comprising a genetic component one desires to introduce into  
the genome of said wheat plant;

(c) identifying or selecting a transformed cell line; and

25 (d) regenerating a transgenic wheat plant therefrom.

19 The method of claim 14 or 18, wherein said DNA comprises an *nptII* gene.

20. The method of claim 18, wherein said *A. Agrobacterium* is *A. tumefaciens* C58






FIG. 1




FIG. 2




FIG. 3




FIG. 4

5/12




FIG. 5

6/12




FIG. 6




FIG. 7




FIG. 8



9/12




FIG. 9





FIG. 10

**PROTOCOL 1**  
**TRANSFORMATION OF IMMATURE EMBRYOS**



**FIG. 11**

PROTOCOL 2  
SUSPENSION CELL TRANSFORMATION

WHEAT CV. MUSTANG  
SUSPENSION CELLS

Subculture 1-3d ↓

INOCULATION WITH *AGROBACTERIUM*

30 min-3 h ↓

CO-CULTURE

2-3 d ↓

DELAY OF SELECTION

1-3 d ↓

SELECTION

40-60 d ↓ Selective agent

TRANSFORMED COLONIES

FIG. 12

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>C12N 15/82, A01H 5/00</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 97/48814</b> <b>(43) International Publication Date:</b> 24 December 1997 (24.12.97)
<b>(21) International Application Number:</b> PCT/US97/10621 <b>(22) International Filing Date:</b> 20 June 1997 (20.06.97) <b>(30) Priority Data:</b> 08/667,188                      21 June 1996 (21.06.96)                      US <b>(71) Applicant:</b> MONSANTO COMPANY [US/US]; 800 North Lindbergh Boulevard, St. Louis, MO 63167 (US). <b>(72) Inventors:</b> CHENG, Ming; 11532 Tivoli Lane, St. Louis, MO 63146 (US). FRY, Joyce, E.; 9839 Gerald Drive, St. Louis, MO 63128 (US). WAN, Yuechun; 893-F Foxsprings, Chesterfield, MO 63017 (US). <b>(74) Agent:</b> KITCHELL, Barbara, S.; Arnold, White & Durkee, P.O. Box 4433, Houston, TX 77210 (US).	<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>  <b>(88) Date of publication of the international search report:</b> 19 March 1998 (19.03.98)	
<b>(54) Title:</b> METHODS FOR THE PRODUCTION OF STABLY-TRANSFORMED, FERTILE WHEAT EMPLOYING <i>AGROBACTERIUM</i> -MEDIATED TRANSFORMATION AND COMPOSITIONS DERIVED THEREFROM		
<b>(57) Abstract</b> <p>Disclosed are processes for producing stably transformed fertile wheat and a system of transforming wheat <i>via Agrobacterium</i>. This invention provides methods for transforming a variety of explants, such as freshly isolated or pre-cultured immature embryos, embryogenic callus and suspension cells. Also disclosed are methods for recovering transgenic plants after transformation within a short period of time, if the explants are regenerable at the time of transformation. Thus, the frequency of somaclonal variation associated with prolonged <i>in vitro</i> culture period is significantly reduced. The transformation frequency using this system is comparable to or better than published methods using other systems, such as microprojectile bombardment.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

# INTERNATIONAL SEARCH REPORT

1st Application No

PCT/US 97/10621

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C12N15/82 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BIOLOGICAL ABSTRACTS, vol. BA91, Philadelphia, PA, US; abstract no. 57286, HESS D ET AL: "TRANSFORMATION EXPERIMENTS BY PIPETTING AGROBACTERIUM INTO THE SPIKELETS OF WHEAT TRITICUM-AESTIVUM L." XP002046366 see abstract & PLANT SCI (LIMERICK) 72 (2). 1990. 233-244. CODEN: PLSCE4 ISSN: 0168-945,	1-3,8,9, 11-13
X	EP 0 709 462 A (MONSANTO CO) 1 May 1996	1-13
Y	see page 3, line 47 - line 52 ---	14-20
	-/-	

☒ Further documents are listed in the continuation of box C

☒ Patent family members are listed in annex

### \* Special categories of cited documents

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*G\* document member of the same patent family

Date of the actual completion of the international search

10 November 1997

Date of mailing of the international search report

04-02-1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel: (+31-70) 340-2040, Tx 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Maddox, A

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/10621

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
Y	MOONEY P A ET AL.: "AGROBACTERIUM TUMEFACIENS-GENE TRANSFER INTO WHEAT TISSUES" PLANT CELL, TISSUE AND ORGAN CULTURE, vol. 25, 1 January 1991, pages 209-218, XP000560763 see the whole document	14-20
X	--- VASIL, V., ET AL.: "Rapid production of transgenic wheat plants by direct bombardment of cultured immature embryos" BIOTECHNOLOGY, vol. 11, December 1993, pages 1553-1558, XP000590888 see the whole document	1-5,7-13
X	--- VASIL, V., ET AL.: "Herbicide resistant fertile transgenic wheat plants obtained by microprojectile bombardment of regenerable embryogenic callus " BIOTECHNOLOGY, vol. 10, June 1992, pages 667-674, XP002046360 see the whole document	3,6
P,X	--- PUKHAL'SKII, V.A., ET AL.: "Genetic transformation of wheat (Triticum aestivum L.) by Agrobacterium tumefaciens" BIOLOGICAL ABSTRACTS DATABASE, DN-99448911, XP002046361 see abstract & GENETIKA, vol. 32, no. 11, 1996, pages 1596-1600,	1,2,8,9, 11-13
A	--- WO 89 12102 A (TEXAS A & M UNIVERSITY SYST) 14 December 1989 see example 8	1-20
A	--- CHENG, M., ET AL.: "Efficient stable transformation of wheat suspension cells mediated by Agrobacterium tumefaciens" IN-VITRO, vol. 32, no. 3 PART2, March 1996, page 105A XP002046362 see abstract P-1170 --- -/--	3,9,17



## INTERNATIONAL SEARCH REPORT

Patent Application No

PCT/US 97/10621

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
A	KORNIENKO, O.V., ET AL.: "Transformation of the wheat <i>Triticum aestivum</i> with a binary vector system of <i>Agrobacterium</i> plasmids" COMMONWEALTH AGRICULTURAL BUREAU DATABASE, DN-931642242, XP002046363 see abstract & (1991) PP. 20-21, 128-129. PUSHCHINO MEETING INFO.: 1 VSESOYUZNYI SIMPOZIUM "NOVYE METODY BIOTEKHNOLOGII RASTENII", PUSHCHINO, 20-22 NOYABRYA, 1991: TEZISY DOKLADOV. SECONDARY SOURCE: REFERATIVNYI ZHURNAL (1992) 6YA313, ---	8,9
A	ISHIDA, Y., ET AL.: "High efficiency transformation of maize ( <i>Zea mays</i> L.) mediated by <i>Agrobacterium tumefaciens</i> " BIOTECHNOLOGY, vol. 14, no. 6, June 1996, pages 745-750, XP002046364 see the whole document ---	14-20
A	EP 0 672 752 A (JAPAN TOBACCO INC) 20 September 1995 see the whole document ---	14-20
A	LANGRIDGE, P., ET AL.: "Transformation of cereals via <i>Agrobacterium</i> and the pollen pathway: a critical assessment" THE PLANT JOURNAL , vol. 2, no. 4, 1992, pages 631-638, XP002046365 see page 632, right-hand column -----	16,20

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/US 97/10621

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0709462 A	01-05-96	US 5631152 A	20-05-97
		CA 2161387 A	27-04-96
-----			
WO 8912102 A	14-12-89	AU 3756889 A	05-01-90
		CN 1042638 A	06-06-90
		DK 285590 A	31-01-91
		EP 0419533 A	03-04-91
		IE 65516 B	01-11-95
		IL 90440 A	16-10-96
		JP 3504800 T	24-10-91
		PT 90709 B	18-07-95
		US 5164310 A	17-11-92
-----			
EP 0672752 A	20-09-95	AU 7546094 A	22-03-95
		CA 2148499 A	09-03-95
		WO 9506722 A	09-03-95
-----			